

COLONOSCOPY (For Patients With Severe Constipation) PREP PACKET

Procedure Date:	Ordering Physician	Procedure Location:
Arrival Time:am/pm	Dr. Ranvir Singh Dr. Stephen Rashbaum	Emory Johns Creek Outpatier
Procedure Time:am/pm	Dr. Nitin Parikh	Northside Forsyth Outpatien
You will be charged \$250 if you no show, cancel/reschedule without 48 hours notice. 770-227-2222		Johns Creek Endoscopy Suite

BOWEL PREP:

- 1. We have forwarded a prescription for MoviPrep to your pharmacy. In addition, purchase Dulcolax Tablets and 1 Bottle of Magnesium Citrate.
- 2. You will need to be on a strict CLEAR **LIQUID DIET ONLY**, the entire day *prior* to your scheduled procedure.
- 3. At **4:00 pm**, on the day *prior* to your scheduled procedure, you will take 2 Dulcolax Tablets as directed.
- 4. At **6:00 pm** on the day *prior* to your scheduled procedure, you will initiate mixing and drinking your first dose of Moviprep.

Take 1 packet of Powder A and 1 packet of Poweder B and place them in the clear 32 oz container supplied by the pharmacy. Mix the powders with 32 oz of a sweet clear liquid. Continue drinking the mixture until gone.

- 5. At **7:00 pm**, you will then consume the full bottle of Magnesium Citrate.
- 6. 5 1/2 hours *prior* to your scheduled procedure am, you will repeat step 4.

- 7. You may continue a clear liquid diet until you initiate step 4. YOU MAY HAVE NOTHING BY MOUTH for 4 hours prior to your procedure.
- 8. NOTE: YOU MAY HAVE NOTHING TO **EAT, SMOKE or CHEW** after midnight. You may only have your clear liquids with your second dose of Moviprep.

DRIVING INSTRUCTIONS:

Please arrange for a responsible adult to drive you to and from the facility. They must plan on staying at the facility during the procedure. Your procedure may be cancelled should you arrive without a driver.

It is best if you can have a responsible adult with you for at least 8-hours following your procedure.

You will not be permitted to drive for 24-hours after the procedure due to anesthesia. You should also refrain from making any important and financial decisions during this time.

** You may want to bring or wear warm socks, as the procedure area tends to be cool.

nt

- -Take your **Heart & Blood Pressure** medications as instructed unless your doctor instructs you otherwise.
- -If you have **Diabetes**, please follow the instructions below:
- *Do not take your diabetes **pills** the morning of the procedure. Feel free to bring them with you to take following the procedure.
- *If you take **Regular Insulin**, do not take it the morning of your scheduled procedure. Please bring it with you to take following your procedure.
- *If you take NPH, or Lantus-type Insulin, take ½ of your regular dose the morning of your scheduled procedure.

-Do NOT take any of the following **Medications:**

*Coumadin-**Stop** 3 Days Prior

*Plavix, Aspirin, Ibuprofen, Iron and Multivitamins-**Stop** 7 Days Prior

Please remove all jewelry prior to your procedure.